

Abstract

■ Full Text

Supplementary Material

PDF (410K)

Contents

Archive

Related material:

PubMed related arts

GO

PubMed articles by:

Björklund, O.

Kahlström, J.

Salmi, P.

Fredholm, B.

PLoS ONE. 2008; 3(12): e3977.

PMCID: PMC2597749

Published online 2008 December 18. doi: 10.1371/journal.pone.0003977.

[Copyright](#) Bjorklund et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Perinatal Caffeine, Acting on Maternal Adenosine A₁ Receptors, Causes Long-Lasting Behavioral Changes in Mouse Offspring

Olga Björklund,^{#*} Johan Kahlström,[#] Peter Salmi,[#] and Bertil B. Fredholm

Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden

Eric Warrant, *Editor*

Lund University, Sweden

[#]Contributed equally.

* E-mail: Olga.Bjorklund/at/gmail.com

Conceived and designed the experiments: PS BBF. Performed the experiments: OB JK PS. Analyzed the data: OB JK PS BBF. Contributed reagents/materials/analysis tools: OB. Wrote the paper: PS BBF.

Received September 21, 2008; Accepted November 14, 2008.

ABSTRACT

TOP

■ ABSTRACT

INTRODUCTION

RESULTS

DISCUSSION

MATERIALS AND
METHODS

SUPPORTING
INFORMATION

REFERENCES

Background

There are lingering concerns about caffeine consumption during pregnancy or the early postnatal period, partly because there may be long-lasting behavioral changes after caffeine exposure early in life.

Methodology/Principal Findings

We show that pregnant wild type (WT) mice given modest doses of caffeine (0.3 g/l in drinking water) gave birth to offspring that as adults exhibited increased locomotor activity in an open field. The offspring also responded to cocaine challenge with greater locomotor activity than mice not perinatally exposed to caffeine. We performed the same behavioral experiments on mice heterozygous for adenosine A₁ receptor gene (A₁RH_z). In these mice signaling via adenosine A₁ receptors is reduced to about the same degree as after modest consumption of caffeine. A₁RH_z mice had a behavioral profile similar to WT mice perinatally exposed to caffeine. Furthermore, it appeared that the mother's genotype, not offspring's, was critical for behavioral changes in adult offspring. Thus, if the mother partially lacked A₁ receptors the offspring displayed more hyperactivity and responded more strongly to cocaine stimulation as adults than did mice of a WT mother, regardless of their genotype. This indicates that long-term behavioral alterations in the offspring result from the

maternal effect of caffeine, and not a direct effect on fetus. WT offspring from WT mother but having a A₁R Hz grandmother preserved higher locomotor response to cocaine.

Conclusions/Significance

We suggest that perinatal caffeine, by acting on adenosine A₁ receptors in the mother, causes long-lasting behavioral changes in the offspring that even manifest themselves in the second generation.

TOP

ABSTRACT

■ INTRODUCTION

RESULTS

DISCUSSION

MATERIALS AND
METHODS

SUPPORTING
INFORMATION

REFERENCES

INTRODUCTION

Caffeine (1,3,7-trimethylxanthine) is a widely consumed psychoactive substance that is readily available through several dietary products (coffee, tea, cocoa beverages and chocolate bars). The total worldwide consumption of caffeine (irrespective of source) has been estimated to approximately 70 to 76 mg/person/day. Interestingly, the levels of caffeine intake in countries such as Sweden and Finland reach more than 400 mg/person/day [1]. Although health consequences of ordinary caffeine consumption are probably minor there are concerns about caffeine intake during pregnancy and lactation. It is notable that in contrast to alcohol and tobacco consumption during pregnancy, approximately 70% of expectant mothers continue to drink beverages containing caffeine at normal or near normal rate [2], [3].

Human and animal studies have shown that high caffeine intake represents a risk for adverse pregnancy outcomes and teratological consequences in offspring [4], [5]. There are many animal studies on the effect of caffeine intake by dams, and often rather high doses of this substance (>50 mg/kg) have been studied. It is therefore important to note that the behavioral effects of caffeine are characterized by a biphasic dose-effect relationship. At low to moderate doses (50 to 300 mg, i.e. 1 to 3 cups of coffee), caffeine induces a central stimulation in humans, eliciting feelings of wellbeing, alertness, energy and ability to concentrate. In contrast, the subjective effects induced by caffeine at higher doses (300 to 800 mg) are characterized by negative feelings such as anxiety, nervousness and insomnia, a condition sometimes referred to as “caffeinism” [6]. In laboratory animals the behavioural effects of caffeine are also biphasic [7]. For example, low doses (<25 mg/kg) of caffeine are similar to psychomotor stimulants such as cocaine and amphetamine, whereas at higher doses caffeine has effects that are similar to a diverse set of other agents such as benzodiazepine-inverse agonists and phencyclidine (PCP) [8].

In low doses, which are the most relevant to human use, caffeine effects are exerted by antagonizing brain adenosine A₁ and A_{2A} receptors with secondary effects on dopaminergic neurotransmission [1]. There is little evidence that these doses produce teratological effects [9]. One concern about early exposure to low or modest doses of caffeine relates to hyperactivity in late adolescence or adulthood [10]. Early exposure to psychostimulant drugs may lead to a phenomenon called “neuronal imprinting” where a drug may have effects that are not necessarily immediate but manifest later in life [11]. Relating to the fact that the rewarding properties of all psychostimulants, including to some extent caffeine [7], are a result of actions of the drugs on the mesolimbic dopamine system [12], early exposure to caffeine might also produce late consequences e.g. in the reaction to other psychoactive drugs.

It is often tacitly assumed that the reason that psychoactive drugs have long-term behavioral consequences is due to the affected fetal brain. However, it is clearly also possible that the drug affects maternal physiology or behavior in such a way that there are long-term consequences in the offspring. It will always be difficult to discriminate between these two possibilities when only drug administration is used. We therefore wanted to see if aspects of the effects of caffeine could be mimicked by genetic targeting of one of the adenosine receptors, since this might allow a separation into maternal or filial effects.

The present study was designed to further assess the behavioural status, including motor functions

and psychomotor activation of adult animals whose mothers were exposed to caffeine during pregnancy and lactation and their response to the psychostimulant cocaine. The rationale behind testing the response to another psychoactive stimulus in these animals is that caffeine, itself a motor stimulant, might be expected to change the motor activity in response to cocaine. Some neurochemical measurements were also made. We found that some effects of perinatal exposure to caffeine were mimicked in mice heterozygous for adenosine A₁ receptors, which have half the normal number of such receptors. This is relevant because regular human consumption of caffeine leads to the blockade of about half of the body's A₁ receptors.

TOP

ABSTRACT

INTRODUCTION

■ RESULTS

DISCUSSION

MATERIALS AND METHODS

SUPPORTING INFORMATION

REFERENCES

RESULTS

Perinatal caffeine exposure

Adult WT mice, 8–10 weeks of age, were perinatally exposed to 0.3 g/l caffeine in the drinking water given to the dams from GD1 to PND21. The dose of caffeine given produced blood levels in dams comparable to those obtained in humans after consumption of 3–4 cups of coffee [13]. Together with offspring of untreated WT dams, caffeine pre-treated male and female mice were tested for motor balance and coordination on the rotarod on Day 1 of the evaluation protocol (Figure 1). There were no major differences in either sex between untreated group and mice given perinatal caffeine in terms of their ability to remain on the rotarod (Figure S1. A,B of the supplemental data).



Figure 1
Experimental design.

Two periods of habituation to the open field were performed on Day 3 and the spontaneous activity was evaluated in all groups tested. As shown in Figure 2A caffeine pre-treated adult WT females displayed increased (121.7 ± 2.1) spontaneous motor activity compared to the WT untreated females (113.3 ± 1.2) ($p = 0.0008$, $t = 3.68$, $df = 35$ Student's *t* test). Adult WT males pre-treated with caffeine also had significantly increased horizontal activity during the 45 min of testing compared to the WT untreated group as already published [14]. In agreement with these results, male and female offspring from our other recent study, where the wild type mothers were treated with the same concentration of caffeine (0.3 g/l) but from GD7 to PND7, also displayed higher locomotor motor activity as adults [14]. This implies that even a narrower time window of caffeine exposure during development might be sufficient in inducing life-long consequences.





Figure 2
Spontaneous horizontal activity measurements after caffeine exposure.

The WT male mice perinatally exposed to caffeine displayed a more pronounced increase in locomotor activity than controls after the cocaine injection [14]. Caffeine pre-treated female mice were also characterized by a higher response to cocaine stimulation than the controls during the 90 min recording session (Figure 2B) (Student's *t* test for H_a: WT H₂O 46.5 ± 3.5 , $n = 8$; WT Caff 73.8 ± 4.7 , $n = 7$, $p = 0.0004$, $t = 4.75$, $df = 13$). This observation was similar to our previous report on the higher effect of amphetamine stimulation in caffeine pre-treated female offspring [14].

Response to cocaine stimulation in A₁R KO and A_{2A}R KO mice

It is known that behavioural effects of 15 mg/kg caffeine can be largely accounted for by blockade of adenosine receptors and that A_1 and A_{2A} receptors are particularly important [1]. We examined if complete elimination of A_1 R would mimic the effect of caffeine. As seen in [Figure 3A,B](#) cocaine injection (10 mg/kg) induced a higher locomotor activity in the A_1 R KO adult male mice than in the wild types (accumulated beam breaks for 90 min following injection for Ha, WT: 39.5 ± 2.6 , $n = 10$, A_1 R KO: 63.9 ± 14.4 , $n = 3$, $p = 0.008$, $df = 11$ Student's t test). This was partially due to a higher basal activity ([Figure 3A](#)). The difference in cocaine response was, if anything, more pronounced in female mice from A_1 R KO genotype than in the WT group of the same sex (Ha, accumulated beam breaks from 46–135 min after cocaine injection: WT 51.6 ± 3.7 , $n = 12$, A_1 R KO: 71.6 ± 6.9 , $n = 4$, $p = 0.009$, $df = 14$ Student's t test), in part as a consequence of a higher basal activity ([Figure 3B](#)).




Whereas A_1 R KO mice appear to be more responsive to cocaine than wild type animals the opposite is known to be true for A_{2A} R KO mice since several previous reports, from our and other groups, have demonstrated decreased response to cocaine or amphetamine in animals that lack A_{2A} receptor gene [14], [15].

The obtained results, higher response of A_1 R KO animals, as well as impaired response in A_{2A} R KO mice to cocaine stimulus reported in the literature, prompted us to continue the work involving adenosine A_1 receptor. Since normal doses of caffeine block half the A_1 receptors in the body [16], we used mice heterozygous for adenosine A_1 receptors as comparison to the WT perinatally caffeine treated mice.

A_1 R Hz mice


Autoradiography with the selective A_1 R antagonist [3 H]DPCPX determined that half of the number of A_1 R present in A_1 R Hz mice than in the WT when the area of nucleus accumbens was examined ([Figure 4](#)) and striatum (not shown).



When A_1 R Hz animals were tested for motor coordination on the rotarod, decreased fall latency was observed in A_1 R Hz males, but no changes in the ability to remain on the rotating rod were found in A_1 R Hz females compared to the wild types ([Figure S1](#), A,B of the supplemental data, males: WT H_2O 124.2 ± 7.4 , $n = 21$ and A_1 R Hz 94.0 ± 9.8 , $n = 8$, $p = 0.03$, $t = 2.2$, $df = 27$ Student's t test).

When tested for the spontaneous activity A_1 R Hz male group did not show a significant increase in locomotion compared to the WT untreated group ([Figure 5A](#)) (45 min of habituation 1, Ha: WT H_2O 110.7 ± 1.0 , $n = 17$; A_1 R Hz 113.6 ± 1.6 , $n = 7$, $p = 0.15$, $t = 1.51$, $df = 22$, Student's t test). A statistically significant increase in locomotion was, however, displayed by female A_1 R Hz group compared to their WT controls ([Figure 5B](#)). Student's t test was run on the score of accumulated beam breaks for

the 45 min period of the first habituation, WT H₂O 113.3±1.2, n = 25 and A₁R Hz 124.1±4.1, n = 8, p = 0.0018, t = 3.4, df = 31.




The enhanced response to cocaine challenge found in WT caffeine pre-treated male groups was also observed in the mice that were heterozygous for the A₁ receptor gene (Figure 5C) (accumulated beam breaks for 90 min recording of Ha: WT H₂O 42.4±3.7, n = 11; A₁R Hz 63.3±1.1, n = 3, p = 0.02, t = 2.8, df = 12 Student's t test). A₁R Hz female mice were also characterized by a higher response to cocaine stimulation than the controls during the 90 min recording session (Figure 5D) (Student's t test, WT H₂O 46.5±3.5, n = 8, A₁R Hz 63.8±4.7, n = 4, p = 0.02, df = 10).

Both adenosine A₁ receptor heterozygous mice and WT mice exposed to perinatal caffeine showed the tendency towards increased locomotion and decreased habituation (Figure 5). Thus, the behavioral profile of A₁R Hz mice which were born by A₁R Hz mothers was basically similar to that of mice exposed to perinatal caffeine. However, part of the increase in response to cocaine in the mice perinatally treated with caffeine or the A₁R Hz group could be due to a higher basal activity since those two groups tended to habituate less completely to the open field arena than the control animals (Figure 5C,D).

Role of maternal genotype

The fact that some of the characteristics of perinatal caffeine exposure could be mimicked by partial deletion of a gene opens the possibility of examining whether its effects in the mother or in the offspring during the perinatal period that determine the phenotype of the offspring in adulthood. We therefore compared the behavior of wild type pups and pups heterozygous for adenosine A₁ receptors born to and raised by mothers heterozygous for adenosine A₁ receptors, with that of pups heterozygous for adenosine A₁ receptors born to and raised by wild type mothers. The latter was achieved by mating WT dams with A₁R KO male mice.

As shown in Figure 6A, there was a statistically significant increase in locomotor activity in male mice (regardless of their genotype) to the response to cocaine injection only when born to a mother that partially lacked adenosine A₁ receptors (significant interaction being mother's genotype p < 0.0001, F_(1,31) = 23.8, Two Way ANOVA with factors offspring and mother's genotype). The same phenomenon was also found in the female offspring of the A₁R Hz mothers after the cocaine stimulation (interaction mother's genotype p = 0.0014, F_(1,27) = 12.6, Two Way ANOVA) (Figure 6B).



Furthermore, as exemplified by female mice in [Figure 2B](#) and [5B](#), the habituation profile of adenosine A₁ receptor heterozygous offspring was similar to that of mice exposed to perinatal caffeine. Thus, the hyperactivity profile in offspring seems to be strongly dependent on whether the mother was heterozygous for adenosine A₁ receptors or not. We have also preformed the statistical analysis of all our data taking into consideration the litter-related issues and observed that the significant results remained.

For additional evaluation of the effect of knocking out one copy of the mother's A₁R gene on the second generation of the offspring we have examined the behaviour of the WT male mice whose mothers were WT but grandmothers were either WT or A₁R Hz mice. After stimulation with cocaine we could still observe modifications caused by the absence of A₁R on the offsprings' reaction to the psychostimulant ([Figure 6C](#)) (accumulated beam breaks from 46–135 min in Ha: WT grandmother WT mother WT offspring: 39.8±3.3, n=9; A₁R Hz grandmother WT mother WT offspring: 51.7±3.8, n=4, p=0.02, t=2.1, df=11 Student's t-test).

Expression of immediate early genes and dopamine receptor subtypes

Some attempts were made to find a neurochemical correlate to the behavioral changes. The dose of cocaine used was too low to induce any significant change in the expression of either NGFI-A mRNA or c-fos in caudate-putamen, nucleus accumbens core, nucleus accumbens shell and medial prefrontal cortex in the WT mice pre-treated with caffeine and the WT controls of both sexes. Also the tyrosine hydroxylase in substantia nigra, [Table 1](#) and [³H]mazindol binding in caudate-putamen (fmol/mg protein, offspring from the WT mothers: 418.8±26.3, n=11; offspring from the A₁R Hz mother: 362.7±21.0, n=26, p=0.07, t=1.5, df=35 Student's t-test), failed to demonstrate any significant difference. We also found no changes in the binding of dopamine D₁ (with SCH23 390) or D₂ (with raclopride) receptor ligands nor in expression of preprotachykinin, prodynorphin, prosomatostatin mRNA levels in nucleus accumbens or caudate-putamen ([Table 1](#)), but there was a significant increase in preproenkephalin mRNA in the offspring born to A₁R Hz mother in the caudate-putamen ([Table 1](#)).

Table 1

Expression of mRNA levels in cocaine-treated offspring to adenosine A₁R Hz or WT mother.

[TOP](#)

[ABSTRACT](#)

[INTRODUCTION](#)

[RESULTS](#)

■ [DISCUSSION](#)

[MATERIALS AND METHODS](#)

[SUPPORTING INFORMATION](#)

[REFERENCES](#)

DISCUSSION

We have confirmed that early exposure to caffeine leads to altered motor behavior, including enhanced responses to cocaine [9], [14], [17], [18], [19]. It is recognized that low doses of cocaine and caffeine produce additive effects [20], [21]. However, in the present experiments caffeine was administered long before cocaine and another explanation than the direct drug interaction must be sought. In most, but not all of these earlier studies, perinatal caffeine in doses similar to those we used caused a slight hyperlocomotion of the offspring even in the absence of a drug challenge. Caffeine and cocaine are both behavioral stimulants, but use somewhat different mechanisms to produce their effects [22], [23]. The mechanisms underlying the behavioral features associated with caffeine, and psycho-stimulants in general, are believed to be related to the activation of dopamine (DA) receptors in the mesolimbic dopamine system (notably in the nucleus accumbens). Whereas cocaine and amphetamine primarily stimulate D₁ receptor, caffeine largely acts by enhancing D₂ receptor pathways [23], [24]. The effects of caffeine is believed to be indirect and due to blockade of adenosine A_{2A} receptors that are co-localized with D₂ receptors that have opposite actions [1].

We found in our recently published study that an even shorter period of maternal caffeine intake (two weeks prenatally and one week postnatally) was sufficient to produce long-lasting behavioral

changes in the offspring [14]. This suggested that the antenatal period is particularly important as also attested by the previous cross-fostering experiments [25]. We cannot rule out that also shorter periods of caffeine treatment before the pregnancy could have effects, but most data on humans suggest a complete return of physiological functions after cessation of caffeine use, for reference see [1]. In female mice, we found no evidence that long term oral caffeine intake in the doses given here alters overall activity, or the location of activity (central vs. peripheral), which may be somewhat related to anxiety following a period of caffeine exposure (Eriksson, Yang, Salmi and Fredholm, unpublished data). Furthermore, we did not observe any gross alteration in maternal behavior in the caffeine treated dams. We therefore favour the hypothesis that maternal caffeine use during a critical period of fetal development is the most important.

In practically all previous studies on perinatal caffeine effects, the focus in the discussion has been on the fetus. Indeed, caffeine readily crosses the placenta without metabolism and partially enters breast milk [26]. The human fetus and newborn infant is exposed to caffeine for a prolonged period of its early life as liver enzymes which metabolize caffeine are not present until eight months of age [26]. However, it is clearly also possible that the relevant action is on the mother. Previous cross-fostering experiments have demonstrated that the effect of a single caffeine dose on the subsequent behavior could not be ascribed to changes in mother's behavior or lactational efficiency in mice [25]. It could, however, be due to an altered uterine environment [27], [28], [29]. It has been shown that a single, very high dose of caffeine (120 mg/kg) could reduce blood flow to the uterus and decidua [30]. This may cause changes in fetal oxygen supply and/or changes in maternal blood composition, but the effects of such high doses are very different from what is observed with the present dosing. However, to completely differentiate between maternal and fetal actions in this period will be virtually impossible when examining drug administration, which by necessity must be through the mother. This is why we tried to find a genetic model that could mimic some aspects of exposure to caffeine.

The present data show that several features of perinatal caffeine administration, notably the enhanced response to cocaine, can be mimicked by deletions of the A_1 receptor, a known target of caffeine, whereas it was known since before that deletion of both copies of the A_{2A} receptor had the opposite effect [15]. We also found that the deletion of only one of the copies of the A_1 receptor gene, resulting in approximately half the number of receptors in cortex and hippocampus [16] but also in regions of more direct relevance here (present data), enhanced the response to cocaine. By contrast, deletion of one of the copies of the A_{2A} gene reduces the response to cocaine (Jiang-Fan Chen, personal communication) albeit not to the same extent as removal of both copies [15]. It is important to note that the A_1 heterozygous mouse still responds to adenosine, but approximately twice as much of the agonist is needed for the same response [16], [31]. Such a parallel shift of the dose-response is also achieved by caffeine at concentrations close to the K_d for the antagonist (10–30 μ M), concentrations that are attained by the doses of caffeine used in the present experiments.

It is also noteworthy that the effect of caffeine on the behavior of the offspring was (at least partially) mirrored by the A_1 receptor deletion, whereas the stimulatory responses of caffeine are generally dependent on the A_{2A} receptor [32], [33] although A_1 receptors contribute [34], [35], [36], [37]. This suggests that the effect is not simply due to psychostimulant actions where A_{2A} receptors are very important, but that some other features must be involved. It is probably relevant that we have previously demonstrated that the ability of a low dose of caffeine to cause reinstatement of cocaine-seeking behavior could be mimicked by A_1 -, but not by A_{2A} receptor antagonists [38], even though A_{2A} antagonists per se tend to produce at least as large increases in locomotion that do A_1 R antagonists [35], [36], [39].

Although a link between the perinatal caffeine use and dopaminergic mechanisms is possible the above data and considerations suggest that the relationship is not simple. Whereas the blockade of A_{2A} receptors has been linked to psychostimulation [32], [33], blockade of A_1 receptors may instead

be related to enhanced glutamatergic transmission [1], [16]. Our examination of various neurochemical parameters did not provide any direct evidence for a major disruption of the dopaminergic pathways involved in psychostimulation. We did find an elevation of preproenkephalinA mRNA levels in caudate-putamen. Enkephalin-expressing neurons project to globus pallidus, a brain region that has been demonstrated to be smaller in children with attention-deficit hyperactivity disorder [40]. The potential association between perinatal exposure to caffeine and ADHD in later life needs to be further evaluated. The precise mechanisms underlying the behavioural enhancement in the mouse offspring therefore remains to be explained. It was also not the major focus of this study.

Instead the aim was to use the genetic model to try to elucidate whether the enhancement of cocaine responses in the offspring was due to a primary effect on the mother or the fetus/pup. The approach we used with a genetic study makes it possible to investigate this aspect, which would not be possible using drugs (since it is impossible to administer psychoactive drugs only to the dam and not to the fetus, and vice versa). The results very clearly indicate the first possibility. Thus, if the mother partially lacked A₁ receptors, the offspring displayed hyperactivity during habituations and responded more strongly to cocaine as adults, regardless of their own genotype. Also, the offspring of wild type mothers showed no behavioral changes, even if they themselves lacked A₁ receptors (i.e. their fathers lacked A₁ receptors). The apparent lack of importance of the fetal brain A₁ receptors is supported by other evidence. Even if adenosine A₁ receptors are present, albeit sparse, in the embryonic brain [41], these receptors appear to be poorly coupled to G proteins [42]. A maternal effect of caffeine related to adenosine A₁ receptor signalling was also shown in a recent study where adenylyl cyclase inhibition by an adenosine A₁ receptor agonist was decreased only in the mother but not the fetal brain [43].

What changes that occur specifically in the pregnant mice that brings about the long-lasting behavioral effects we demonstrate here in the offspring is not known, and will need consideration in future studies. Our experiments performed on the second generation of mice whose grandmother was A₁R Hz but mother a WT still showed increased response to stimulation with cocaine compared to the WT mice whose both mother and grandmother were wild types. This indicates that long-term behavioral alterations in the offspring may greatly depend on a maternal effect of caffeine and not a direct effect in the fetus, and that some epigenetic effect may be involved.

Epigenetic change to the genome (e.g. in DNA methylation) not only determine the phenotype of the offspring but can sometimes be passed on to the second generation. These processes of transgenerational passage of changes in genomic DNA methylation can occur in both female and male lineage, as the transmission is only via the gametes and can equally apply to sperm and ova [44]. Therefore, epigenetical transmission from father or mother could have the effects on the developmental responses in the offspring. In our study, we have controlled for the mother's and grandmother's genotype in A₁R Hz offspring but not for the father's or grandfather's genotype. There were two reasons for not considering the paternal genotype. Firstly we wanted to relate our findings to the exposure of mouse dams to caffeine. Clearly paternal effects can not be very important here. Secondly we noted in a separate study that life-long exposure (including perinatal exposure) to caffeine did not cause behavioral changes in the male but did in female mice (Salmi P, Fredholm BB unpublished data).

In summary, we found that adenosine A₁ receptor heterozygous offspring had a behavioral profile of hyperactivity quite similar to normal mice exposed perinatally to caffeine. Furthermore, it appeared that the genotype of the dam, not the offspring, was critical for behavioral changes in adult offspring. We hypothesize that perinatal caffeine, by acting on adenosine A₁ receptors in the mother, appears to be responsible for the long-lasting behavioral changes in the offspring. It should be remembered, however, that at this point it is not possible to conclude whether the long-lasting behavioral changes observed here are detrimental or beneficial.

[TOP](#)[ABSTRACT](#)[INTRODUCTION](#)[RESULTS](#)[DISCUSSION](#)[MATERIALS AND METHODS](#)[SUPPORTING INFORMATION](#)[REFERENCES](#)

MATERIALS AND METHODS

Animals

Three types of mice were used: *adenosine A₁ receptor knock-out (A₁R KO)* [16], *mouse heterozygous for the adenosine A₁ receptor (A₁R Hz)* and *C57BL/6 mice* (wild type, WT). For generating the A₁R KO mice, the second coding exon of the mouse adenosine A₁ receptor gene was inactivated in mouse E14.1 embryonic stem cells. 129/OlaHsd/C57Bl/6 hybrid mice were generated. These animals were backcrossed for at least 10 generations with C57Bl/6 to achieve practically congenic A₁R KO mice. Animals were bred at the Department of Physiology and Pharmacology, Karolinska Institutet and housed at a constant room temperature (22°C; 12 h light/dark cycle, lights on at 6 a.m) with *ad libitum* access to food and water. To generate the A₁R KO and A₁R Hz mice heterozygote matings were usually performed, except where otherwise stated. Mice were genotyped by PCR. All experiments were approved by the Local Committee on Ethics of Animal Experimentation, Stockholm, Sweden.

Chemicals

Caffeine (anhydrous; Sigma Chemical Co., St. Louis, MO) was administered through the animals' drinking water (0.3 g/l). Cocaine hydrochloride (Apoteksbolaget AB, Sweden) was dissolved in 0.9% NaCl (10 mg/ml) and injected intra-peritoneally (i.p.) in a dose of 10 mg/kg. Saline vehicle served as controls.

Exposure to caffeine

The experimental design of caffeine exposure is shown in [Fig. 1](#). Pregnant WT females, housed in individual cages, were administered caffeine through their drinking water. The first set of animals received 0.3 g/l caffeine and the second was given tap water only, and served as control. The exposure time ranged from gestational day (GD) 1 to day 21 of lactation, i.e. postnatal day (PND) 21. That is why the term “perinatal” is here used to denote the period of pregnancy and the 21 days after birth. The offspring was analyzed with behavioral tests at two months of age.

Behavioral evaluation. All behavioral tests were performed during the light period between 8 a.m. and 4 p.m. The animals were separated into groups based on exposure (caffeine, water, A₁R Hz from A₁R Hz mothers) and stimulation (cocaine or saline). Males and females were analyzed separately (n = 6–25 mice in each exposure group, selected randomly from 2–4 independent litters). The motor activity in an open field arena was analyzed at adult age with an experimental protocol that spanned over 4 days ([Fig. 1](#)). The rotarod test, including the training session, was performed during the first day. Mice were allowed to rest the next day, and on the third day two open field habituation sessions were performed. The challenge with cocaine, preceded by another habituation session, was done on the fourth day. For this purpose the mice from each exposure group were divided into two sets, one stimulated with cocaine, the other receiving saline. A₁R KO and their WT littermates were also analysed according to the 4 day protocol explained above.

Regarding adenosine A₁ receptor heterozygous mice, and their wild type littermates, four different groups of offspring were tested: 1) wild type mice born to an adenosine A₁ receptor heterozygous mother, 2) adenosine A₁ receptor heterozygous mice born to an adenosine A₁ receptor heterozygous mother, 3) adenosine A₁ receptor heterozygous mice born to a wild-type mother, and 4) wild type mice born to a wild type mother. Behavioral tests were performed when offspring were at least 8 week of age. A₁R Hz mice were separated into groups described above and challenged with cocaine.

For the evaluation of the effect of knocking out one copy of the A₁R gene on the second generation

of the offspring we have examined the behaviour of the WT male mice whose mothers were wild type but grandmothers were either WT or A₁R Hz mice. These mice underwent the same behavioural protocol as described in above ([Fig. 1](#)).

Open field model

The effects of caffeine on motor functions were analyzed by recording the locomotor activity in a square open field arena (500×500×225 mm), enclosed in a solid and sound-attenuating box (Kungsbacka Mät och Reglerteknik AB, Fjärås, Sweden). The open field arena was equipped with two rows of photocells sensitive to infrared light, each row having 16 photocells per side. The space between the photocells was 31 mm and the outermost was placed 17.5 mm from the wall. The number of photocell interruptions was collected by a computer and the following variables were recorded and analysed: horizontal activity (Ha, total number of beam breakings), locomotion (L, interruptions of photocells in the lower rows when there is a new beam broken, i.e. the animal has made an actual transfer) and rearing activity (Ra, all interruptions of photocells in the upper rows). This equipment does not allow recording small movements, e.g. tremor, reflexes and tail movements. The data were subjected to a square root transformation (sqrt) before statistical analysis.

Prior to the recording, all animals were allowed a period of 30–45 min in behavioral the testing room. At 2 months of age the mice were analyzed with an experimental protocol that spanned 2 days. Two habituations (45 min each) were performed on day 1, separated by 2 h period. Directly after the first habituation (habituation 3) on the second day the mice were injected with cocaine (10 mg/kg) and their locomotor activity was recorded for 90 min. A₁ receptor heterozygous mice were separated into groups described in 2.3. and challenged with cocaine.

Rotarod

Potential effect of the perinatal caffeine exposure on the cerebellum was tested by Rotarod test at 2 months of age (LSi Leticia Scientific Instruments, Debiomed, Cornella, Spain). Mice were initially trained to remain on the rotating drum at a constant speed of 4 r.p.m. (revolutions per minute) for 3 min. During the three trials the rotarod accelerated from 4 to 40 r.p.m. over a 5 min period. Two mice were tested simultaneously, and every animal was involved in three consecutive trials, each separated by a 30 min resting period. Fall latencies were recorded. Analysis was performed using the best value obtained in 3 consecutive sessions for each animal.

Tissue preparation

After the experiments (approximately 2 h after cocaine or vehicle injection) mice were anaesthetized with CO₂ and sacrificed by decapitation. The brains were dissected out, frozen on dry ice and stored at –80°C. Frozen brains were cut by cryostat into 14 µm coronal sagittal sections, thaw-mounted on poly-L-lysine coated slides as previously described [[45](#)], [[46](#)] and stored at –20°C.

In situ hybridization and receptor autoradiography

The immediate early genes NGFI-A and c-fos were examined by in situ hybridisation on coronal sections (14 µm). Expression of preprotachykinin, prodynorphin, prosomatostatin, preproenkephalin and tyrosine hydroxylase mRNA was also measured [[39](#)], [[47](#)]. Binding studies included dopamine D₁ receptor binding examined by 0.2, 0.4, 0.6, 1 and 2 nM [³H]SCH23390 ((R)-(+)-7-chloro-8-hydroxy-3-methyl-1-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine, DuPont NEN, Stevenage, UK), D₂ receptor binding with 1, 2, 3, 5 and 10 nM [³H]raclopride (DuPont NEN, Stevenage, UK), respectively as described before [[48](#)] and 10 nM [³H]mazindol binding [[49](#)]. A₁R binding was evaluated by incubating the brain section from WT and A₁R Hz animals with increasing (0.2–10 nM) concentrations of A₁R antagonist [³H]1,3-dipropyl-8-cyclopentylxanthine ([³H]DPCPX) as described

in [50].

Films were developed after 4 weeks of exposure. The autoradiographic films were digitized using a CCD camera (Sierra Scientific Sunnyvale, CA, USA), and optical densities were converted to fmol/mg tissue using a Kodak density wedge and the MCID M5 system (Imaging Research, St. Catharines, Canada). Specific binding was obtained by subtracting non-specific binding from total binding.

Statistical analysis

Data were analyzed using GraphPad Prism 4 (GraphPad Software Inc., San Diego, USA). To determine statistical significance, open field data were first normalized with a square root transformation. Accumulated counts (horizontal activity and rearing) and rotarod data (fall latencies) were analysed using Student's t test. In some cases Two Way ANOVA was used, as indicated. Differences were considered statistically significant at $p \leq 0.05$. Data are presented as means \pm S.E.M.

TOP

SUPPORTING INFORMATION

Figure S1

Rotarod measurements. Analysis of the rotarod performance in adult WT and A₁R Hz male (A) and female (B) mice that received perinatally 0.3 g/l caffeine or tap water. Fall latencies (mean \pm S.E.M.) were analyzed by Student's t test (males WT H₂O: n = 21, WT Caff: n = 7, A₁R Hz: n = 8; females WT H₂O: n = 22, WT Caff: n = 13, A₁R Hz: n = 8).

(3.52 MB TIF)

[Click here for additional data file.](#) ^(3.3M)

ACKNOWLEDGMENTS

We thank Dr. Jiang-Fan Chen for critical reading of the manuscript and Karin Lindström for help with the in situ and autoradiography experiments.

FOOTNOTES

Competing Interests: The authors have declared that no competing interests exist.

Funding: This work was supported by Karolinska Institutet, the Swedish Science Council (project No. 2553), European Commission (LSHM-CT2005-518189) and NIH (1R01 NS048995-01A1). The funding agencies do not take any responsibility for the contents of the article.

TOP

ABSTRACT

INTRODUCTION

RESULTS

DISCUSSION

MATERIALS AND
METHODS

SUPPORTING
INFORMATION

■ REFERENCES

REFERENCES

1. Fredholm, BB; Battig, K; Holmen, J; Nehlig, A; Zvartau, EE. Actions of caffeine in the brain with special reference to factors that contribute to its widespread use. *Pharmacol Rev.* 1999;51:83–133. [[PubMed](#)]
2. Olsen, J; Overvad, K; Frische, G. Coffee consumption, birthweight, and reproductive failures. *Epidemiology.* 1991;2:370–374. [[PubMed](#)]
3. Bracken, MB; Triche, EW; Belanger, K; Hellenbrand, K; Leaderer, BP. Association of maternal caffeine consumption with decrements in fetal growth. *Am J Epidemiol.* 2003;157:456–466. [[PubMed](#)]
4. Klebanoff, MA; Levine, RJ; DerSimonian, R; Clemens, JD; Wilkins, DG. Maternal serum paraxanthine, a caffeine metabolite, and the risk of spontaneous abortion. *N Engl J Med.* 1999;341:1639–1644. [[PubMed](#)]

- Cnattingius, S; Signorello, LB; Anneren, G; Clausson, B; Ekbom, A, et al. Caffeine intake and the risk of first-trimester spontaneous abortion. *N Engl J Med.* 2000;343:1839–1845. [[PubMed](#)]
6. Nehlig, A; Boyet, S. Dose-response study of caffeine effects on cerebral functional activity with a specific focus on dependence. *Brain Res.* 2000;858:71–77. [[PubMed](#)]
 7. Daly, JW; Fredholm, BB. Caffeine—an atypical drug of dependence. *Drug Alcohol Depend.* 1998;51:199–206. [[PubMed](#)]
 8. Mumford, GK; Holtzman, SG. Qualitative differences in the discriminative stimulus effects of low and high doses of caffeine in the rat. *J Pharmacol Exp Ther.* 1991;258:857–865. [[PubMed](#)]
 9. Nehlig, A; Debry, G. Potential teratogenic and neurodevelopmental consequences of coffee and caffeine exposure: a review on human and animal data. *Neurotoxicol Teratol.* 1994;16:531–543. [[PubMed](#)]
 10. Linnet, KM; Dalsgaard, S; Obel, C; Wisborg, K; Henriksen, TB, et al. Maternal lifestyle factors in pregnancy risk of attention deficit hyperactivity disorder and associated behaviors: review of the current evidence. *Am J Psychiatry.* 2003;160:1028–1040. [[PubMed](#)]
 11. Andersen, SL. Stimulants and the developing brain. *Trends Pharmacol Sci.* 2005;26:237–243. [[PubMed](#)]
 12. Feldman, RS; Meyer, JS; Quenzer, LF. Principles of Neuropharmacology. *Principles of Neuropharmacology.* 1997:612.
 13. Bona, E; Aden, U; Fredholm, BB; Hagberg, H. The effect of long term caffeine treatment on hypoxic-ischemic brain damage in the neonate. *Pediatr Res.* 1995;38:312–318. [[PubMed](#)]
 14. Bjorklund, O; Kahlstrom, J; Salmi, P; Ogren, SO; Vahter, M, et al. The effects of methylmercury on motor activity are sex- and age-dependent, and modulated by genetic deletion of adenosine receptors and caffeine administration. *Toxicology.* 2007;241:119–133. [[PubMed](#)]
 15. Chen, JF; Beilstein, M; Xu, YH; Turner, TJ; Moratalla, R, et al. Selective attenuation of psychostimulant-induced behavioral responses in mice lacking A(2A) adenosine receptors. *Neuroscience.* 2000;97:195–204. [[PubMed](#)]
 16. Johansson, B; Halldner, L; Dunwiddie, TV; Masino, SA; Poelchen, W, et al. Hyperalgesia, anxiety, and decreased hypoxic neuroprotection in mice lacking the adenosine A1 receptor. *Proc Natl Acad Sci U S A.* 2001;98:9407–9412. Epub 2001 Jul 9424. [[PubMed](#)]
 17. Henderson, MG; McConnaughey, MM; McMillen, BA. Long-term consequences of prenatal exposure to cocaine or related drugs: effects on rat brain monoaminergic receptors. *Brain Res Bull.* 1991;26:941–945. [[PubMed](#)]
 18. Nakamoto, T; Roy, G; Gottschalk, SB; Yazdani, M; Rossowska, M. Lasting effects of early chronic caffeine feeding on rats' behavior and brain in later life. *Physiol Behav.* 1991;49:721–727. [[PubMed](#)]
 19. Peruzzi, G; Lombardelli, G; Abbracchio, MP; Coen, E; Cattabeni, F. Perinatal caffeine treatment: behavioral and biochemical effects in rats before weaning. *Neurobehav Toxicol Teratol.* 1985;7:453–460. [[PubMed](#)]
 20. Gauvin, DV; Criado, JR; Moore, KR; Holloway, FA. Potentiation of cocaine's discriminative effects by caffeine: a time-effect analysis. *Pharmacol Biochem Behav.* 1990;36:195–197. [[PubMed](#)]
 21. Bedingfield, JB; King, DA; Holloway, FA. Cocaine and caffeine: conditioned place preference, locomotor activity, and additivity. *Pharmacol Biochem Behav.* 1998;61:291–296. [[PubMed](#)]
 22. Johansson, B; Lindstrom, K; Fredholm, BB. Differences in the regional and cellular localization of c-fos messenger RNA induced by amphetamine, cocaine and caffeine in the rat. *Neuroscience.* 1994;59:837–849. [[PubMed](#)]
 23. Kuzmin, A; Johansson, B; Fredholm, BB; Ogren, SO. Genetic evidence that cocaine and caffeine stimulate locomotion in mice via different mechanisms. *Life Sci.* 2000;66:PL113–118. [[PubMed](#)]
 24. Hummel, M; Unterwald, EM. D1 dopamine receptor: a putative neurochemical and behavioral link to cocaine action. *J Cell Physiol.* 2002;191:17–27. [[PubMed](#)]

25. Ajarem, JS; Brain, PF. Prenatal caffeine exposure modifies behavioural responses in mice. *Behav Pharmacol.* 1993;4:541–544. [[PubMed](#)]
26. Grosso, LM; Bracken, MB. Caffeine metabolism, genetics, and perinatal outcomes: a review of exposure assessment considerations during pregnancy. *Ann Epidemiol.* 2005;15:460–466. [[PubMed](#)]
27. Pollard, I; Claassens, R. Caffeine-mediated effects on reproductive health over two generations in rats. *Reprod Toxicol.* 1992;6:541–545. [[PubMed](#)]
28. Pollard, I; Jabbour, H; Mehrabani, PA. Effects of caffeine administered during pregnancy on fetal development and subsequent function in the adult rat: prolonged effects on a second generation. *J Toxicol Environ Health.* 1987;22:1–15. [[PubMed](#)]
29. Sinton, CM. Preliminary indications that functional effects of fetal caffeine exposure can be expressed in a second generation. *Neurotoxicol Teratol.* 1989;11:357–362. [[PubMed](#)]
30. Kimmel, CA; Kimmel, GL; White, CG; Grafton, TF; Young, JF, et al. Blood flow changes and conceptual development in pregnant rats in response to caffeine. *Fundam Appl Toxicol.* 1984;4:240–247. [[PubMed](#)]
31. Johansson, SM; Yang, JN; Lindgren, E; Fredholm, BB. Eliminating the antilipolytic adenosine A₁ receptor does not lead to compensatory changes in the antilipolytic actions of PGE₂ and nicotinic acid. *Acta Physiol (Oxf).* 2007;190:87–96. [[PubMed](#)]
32. El Yacoubi, M; Ledent, C; Menard, JF; Parmentier, M; Costentin, J, et al. The stimulant effects of caffeine on locomotor behaviour in mice are mediated through its blockade of adenosine A_{2A} receptors. *Br J Pharmacol.* 2000;129:1465–1473. [[PubMed](#)]
33. Ledent, C; Vaugeois, JM; Schiffmann, SN; Pedrazzini, T; El Yacoubi, M, et al. Aggressiveness, hypoalgesia and high blood pressure in mice lacking the adenosine A_{2a} receptor. *Nature.* 1997;388:674–678. [[PubMed](#)]
34. Ferre, S; Ciruela, F; Borycz, J; Solinas, M; Quarta, D, et al. Adenosine A₁–A_{2A} receptor heteromers: new targets for caffeine in the brain. *Front Biosci.* 2008;13:2391–2399. [[PubMed](#)]
35. Kuzmin, A; Johansson, B; Gimenez, L; Ogren, SO; Fredholm, BB. Combination of adenosine A₁ and A_{2A} receptor blocking agents induces caffeine-like locomotor stimulation in mice. *Eur Neuropsychopharmacol.* 2006;16:129–136. [[PubMed](#)]
36. Karcz-Kubicha, M; Antoniou, K; Terasmaa, A; Quarta, D; Solinas, M, et al. Involvement of adenosine A₁ and A_{2A} receptors in the motor effects of caffeine after its acute and chronic administration. *Neuropsychopharmacology.* 2003;28:1281–1291. [[PubMed](#)]
37. Halldner, L; Aden, U; Dahlberg, V; Johansson, B; Ledent, C, et al. The adenosine A₁ receptor contributes to the stimulatory, but not the inhibitory effect of caffeine on locomotion: a study in mice lacking adenosine A₁ and/or A_{2A} receptors. *Neuropharmacology.* 2004;46:1008–1017. [[PubMed](#)]
38. Kuzmin, A; Johansson, B; Zvartau, EE; Fredholm, BB. Caffeine, acting on adenosine A₁ receptors, prevents the extinction of cocaine-seeking behavior in mice. *J Pharmacol Exp Ther.* 1999;290:535–542. [[PubMed](#)]
39. Svenningsson, P; Georgieva, J; Kontny, E; Heilig, M; Fredholm, BB. Involvement of a c-fos-dependent mechanism in caffeine-induced expression of the preprotachykinin A and neurotensin/neuromedin N genes in rat striatum. *Eur J Neurosci.* 1997;9:2135–2141. [[PubMed](#)]
40. Swanson, J; Castellanos, FX; Murias, M; LaHoste, G; Kennedy, J. Cognitive neuroscience of attention deficit hyperactivity disorder and hyperkinetic disorder. *Curr Opin Neurobiol.* 1998;8:263–271. [[PubMed](#)]
41. Rivkees, SA. The ontogeny of cardiac and neural A₁ adenosine receptor expression in rats. *Brain Res Dev Brain Res.* 1995;89:202–213.
42. Åden, U; Herlenius, E; Tang, LQ; Fredholm, BB. Maternal caffeine intake has minor effects on adenosine receptor ontogeny in the rat brain. *Pediatr Res.* 2000;48:177–183. [[PubMed](#)]
- Leon, D; Albasanz, JL; Ruiz, MA; Martin, M. Chronic caffeine or theophylline intake during pregnancy inhibits A₁

43. receptor function in the rat brain. *Neuroscience*. 2005;131:481–489. [[PubMed](#)]
44. Pembrey, ME. Time to take epigenetic inheritance seriously. *Eur J Hum Genet*. 2002;10:669–671. [[PubMed](#)]
45. Halldner, L; Lozza, G; Lindstrom, K; Fredholm, BB. Lack of tolerance to motor stimulant effects of a selective adenosine A(2A) receptor antagonist. *Eur J Pharmacol*. 2000;406:345–354. [[PubMed](#)]
46. Henry, B; Crossman, AR; Brotchie, JM. Characterization of enhanced behavioral responses to L-DOPA following repeated administration in the 6-hydroxydopamine-lesioned rat model of Parkinson's disease. *Exp Neurol*. 1998;151:334–342. [[PubMed](#)]
47. Svenningsson, P; Strom, A; Johansson, B; Fredholm, BB. Increased expression of c-jun, junB, AP-1, and preproenkephalin mRNA in rat striatum following a single injection of caffeine. *J Neurosci*. 1995;15:3583–3593. [[PubMed](#)]
48. Johansson, B; Georgiev, V; Fredholm, BB. Distribution and postnatal ontogeny of adenosine A2A receptors in rat brain: comparison with dopamine receptors. *Neuroscience*. 1997;80:1187–1207. [[PubMed](#)]
49. Javitch, JA; Strittmatter, SM; Snyder, SH. Differential visualization of dopamine and norepinephrine uptake sites in rat brain using [³H]mazindol autoradiography. *J Neurosci*. 1985;5:1513–1521. [[PubMed](#)]
50. Svenningsson, P; Nomikos, GG; Fredholm, BB. The stimulatory action and the development of tolerance to caffeine is associated with alterations in gene expression in specific brain regions. *J Neurosci*. 1999;19:4011–4022. [[PubMed](#)]

Articles from *PLoS ONE* are provided here courtesy of
Public Library of Science

[Write to PMC](#) | [PMC Home](#) | [PubMed](#)
[NCBI](#) | [U.S. National Library of Medicine](#)
[NIH](#) | [Department of Health and Human Services](#)
[Privacy Policy](#) | [Disclaimer](#) | [Freedom of Information Act](#)